

Floral Preference of Bees in a Montane Meadow in Flagstaff, AZ

Evan Levy¹, Anita Cisternas-Fuentes^{2,3}, Tania Jogesh³, Krissa Skogen³.

¹Colorado College Dept. of Organismal Biology & Ecology, ²Northwestern University, ³Chicago Botanic Garden



Introduction

“The relationship between utilized and available resources in the same environment has not been studied in most wild bees” Dalmazzo *et al.* 2015.

For the few bee species studied, known feeding behaviors range from broadly polylectic to oligolectic^{4, 6, 7}. Evaluating floral resources and bee preferences is essential for understanding the interaction between plants and pollinators in a community. Most studies focus on the lifetime foraging of specific bee taxa, or are done by net sweeping^{2, 4, 6}. This study is unique because it utilizes pan traps to analyze the pollen loads of all bees in a montane community at different times of day. Our hypotheses are:

- H1:** Pollen utilization by bees will not represent available floral resources.
- H2:** Bees will be mainly polylectic (i.e. non-specialists).
- H3:** Floral preference will differ between species of bees.
- H4:** Bees will change dietary preference throughout the day.

Methods

- The study was conducted in a montane meadow in Flagstaff, AZ at an elevation of 2100m (fig. 1)
- White, blue and yellow pan traps were set out between July 2nd and July 29th and checked in the morning (7:00-11:00), afternoon (11:00-16:00) and evening (16:00-20:00) (fig. 2).
- Bees were swabbed for pollen using fuschin gel³ (fig. 3 & 4). The gel was melted onto a glass slide. The bees were pinned for identification.
- Slides were examined under a light microscope at 20x magnification. Pollen grains were counted and identified to the lowest possible taxon.
- A floral survey was conducted to assess pollen availability
- A pollen atlas was made for flowering plants using the same technique with flowers as with bees (fig. 5).
- A chi-squared goodness of fit test and the Kruskal Wallis chi-squared tests were used to analyze data. All tests were run in Rstudio 3.2.1.

Results




Fig. 6: Left: Pictures of each morphospecies with corresponding relative abundance of pollen in morning and afternoon. Sample sizes are indicated above each bar. Right: Results from floral survey based on number of flowers (above) and number of flowering plants (below).

Statistics: Morphospecies 1, 2 and 3 all carried significantly different pollen compared to the pollen resources available based on a chi-squared goodness of fit test ($p < 0.001$). Morphospecies 2 and 3 carried more Asteraceae pollen than morphospecies 1 (Kruskal Wallis $\chi^2 = 9.3$, $p < 0.01$). Morphospecies 2 carried more *M. officinalis* pollen in the morning than in the afternoon (Kruskal Wallis $\chi^2 = 5.7$, $p = 0.017$), whereas morphospecies 1 carried marginally more *Penstemon* pollen in the afternoon than in the morning (Kruskal Wallis $\chi^2 = 3.2$, $p = 0.07$).



Figure 2. Example of a pan trap location.



Figure 3. Morphospecies 3 before and after swabbing



Figure 4. Fuschin gel with bee pollen before melting



Figure 1. Study site in Flagstaff, AZ

Results & Discussion

H1: Bee morphospecies 1, 2 and 3 showed floral preferences that were not proportional to pollen availability, suggesting that floral resource utilization does not match resource availability.

H2: Morphospecies 1, 2 & 3 are broadly polylectic, according to Cane (2006), because they fed on 3+ plant families. Morphospecies 8 & 9 didn't have a large enough sample size to determine preference.

H3: Each morphospecies of bee showed a difference in floral preference. This could be due to different species exploiting different niches.

H4: Floral preference changed between morning and afternoon for morphospecies 1 and 2. These bees could be responding to daily fluctuations in pollen availability, based on when pollen is produced by plants and when it is consumed by insects.




Figure 5: Pollen of 6 different species

H. richardsoni, *A. geyeri*, *C. arvensis*, *P. virgatus*, *P. crinita*, *L. wrightii*

Works Cited

- Cane J.H., et al. 2011. Nectar and pollen sugars constituting larval provisions of the alfalfa leaf-cutting bee (*Megachile rotundata*) (Hymenoptera: Apiformes: Megachilidae). *Apidologie* 42:401–408
- Dalmazzo, Milagro, and Favin Gerardo Vossler. 2015. Pollen host selection by a broadly polylectic halictid bee in relation to resource availability. *Arthropod-Plant Interactions* 9:3: 253-262.
- Xearns, C. A., and D. W. Inouye. 1993. Techniques for pollination biologists. University Press of Colorado.
- Machvor, J. S. et al. 2014 "Pollen specialization by solitary bees in an urban landscape." *Urban ecosystems* 17:1: 139-147.
- O'Neill, Kevin M., et al. 2004. Composition of pollen loads of *Megachile rotundata* in relation to flower diversity (Hymenoptera: Megachilidae). *Journal of the Kansas Entomological Society* 77:4: 619-625.
- Somme, Laurent, et al. 2015. Pollen and nectar quality drive the major and minor floral choices of bumble bees. *Apidologie* 46:1: 92-106.
- Vanderplanck, Maryse, et al. 2014. How does pollen chemistry impact development and feeding behaviour of polylectic bees. *PloSone* 9:1: e86209.

Acknowledgements

We'd like to thank the NSF grant DEB 1342673 for funding this project. Many thanks to the Arboretum at Flagstaff, the Merriam Powell Research Station and Northern Arizona University for providing facilities and study sites to make this study possible. Thank you to Krissa Skogen, Jeremie Fant, Tass Kelso and Rick Overson for providing guidance when needed.

